

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### Ethoxycarbonyl functionalized Tröger's base alongside its congener dihydroquinazoline

Kovalenko, Vitaly; Tumanov, Nikolay; Vasiutovich, Kiryl; Leyssens, Tom

*Published in:*  
Chemical Data Collections

*DOI:*  
[10.1016/j.cdc.2020.100339](https://doi.org/10.1016/j.cdc.2020.100339)

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for pulished version (HARVARD):*  
Kovalenko, V, Tumanov, N, Vasiutovich, K & Leyssens, T 2020, 'Ethoxycarbonyl functionalized Tröger's base alongside its congener dihydroquinazoline: A trick with crystallization', *Chemical Data Collections*, vol. 25, 100339. <https://doi.org/10.1016/j.cdc.2020.100339>

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Contents lists available at ScienceDirect

## Chemical Data Collections

journal homepage: [www.elsevier.com/locate/cdc](http://www.elsevier.com/locate/cdc)

## Data Article

## Ethoxycarbonyl functionalized Tröger's base alongside its congener dihydroquinazoline: A trick with crystallization

Vitaly Kovalenko<sup>a,\*</sup>, Nikolay Tumanov<sup>b</sup>, Kiryl Vasiutovich<sup>a</sup>, Tom Leyssens<sup>c</sup><sup>a</sup> Laboratory of Biochemistry and Biotechnology, Department of Natural Sciences, Belarusian State Pedagogical University, 220030 Minsk, Belarus<sup>b</sup> Department of Chemistry, Université de Namur, 5000 Namur, Belgium<sup>c</sup> Institute of Condensed Matter and Nanosciences, Université catholique de Louvain, B-1348 Louvain-La-Neuve, Belgium

## ARTICLE INFO

## Article history:

Received 10 November 2019

Revised 22 December 2019

Accepted 3 January 2020

Available online 14 January 2020

## Keywords:

Tröger's base

Dihydroquinazolines

Reaction intermediate

Crystallization

Separation

## ABSTRACT

The synthesis of ethoxycarbonyl functionalized Tröger's base is accompanied by the formation of a considerable amount of a 3,4-dihydroquinazoline by-product. We found that both compounds can be readily isolated from reaction mixture on a preparative scale by standard crystallization from appropriate solvents.

© 2020 Elsevier B.V. All rights reserved.

## Specifications table

Subject area	Organic chemistry
Compounds	2,8-Bis(ethoxycarbonyl)-6H,12H-5,11-methanodibenzo[b,f][1,5]diazocine; Ethyl 3-[4-(ethoxycarbonyl)phenyl]-3,4-dihydroquinazoline-6-carboxylate
Data category	Spectral, synthesized, crystallographic, melting points
Data acquisition format	NMR, IR, MS, XRD
Data type	Analyzed
Procedure	Chemical transformations, purification
Data accessibility	Provided with this article

## 1. Rationale

Since the first synthesis by Julius Tröger [1], Tröger's bases continue to fascinate chemists due to their unique structural properties and reactivity. They find a growing number of applications in a variety of domains including supramolecular systems, metallo complexes, polymers, studying important aspects of stereochemistry [2–5]. Herein, we focused on the multigram synthesis of the ester-functionalized Tröger's base **1** (Scheme 1) being of interest to us as a promising object for further chiral resolution. This molecule was used previously for the construction of coordination polymers [6] and receptors which selectively bind dicarboxylic acids [7,8].

\* Corresponding author.

E-mail address: [kovalenkovn@rambler.ru](mailto:kovalenkovn@rambler.ru) (V. Kovalenko).





Fig. 1. Molecular structure of compound 1.



Scheme 1. Synthesis of Tröger base 1 and dihydroquinazoline 2.

According to several thorough studies [9–11], condensation of anilines with paraformaldehyde in a trifluoroacetic acid medium is the method of choice to obtain structural analogues of Tröger's base. This process could be complicated by the formation of unreactive 3,4-dihydroquinazoline semi-products which terminate further reaction sequences [12]. This happens, especially, in the case of anilines with electron withdrawing substituents. It is worth to note that dihydroquinazolines themselves form another diverse family of highly potent nitrogen heterocycles [13–15].

We performed the synthesis of Tröger's base 1 following the Sergeyev et al. procedure [10], having obtained a mixture of target compound 1 and by-product dihydroquinazoline 2 in a poor ratio 2:1 (Scheme 1). Variation of the reaction temperature and aniline 3/paraformaldehyde ratio, or longer reaction time did not improve the yield of 1. In the work [11] this difficulty was circumvented by utilizing *ortho* substituted anilines and esters with longer alkyl chain. In an effort to develop a more preparative procedure for 1, we found that compounds 1 and 2 demonstrate different solubility behavior (despite a structural similarity), and that they can be isolated selectively by crystallization from appropriate solvents. Apparently, dihydroquinazoline 2 fell out from toluene in a yield close to quantitative, while Tröger's base 1 remains soluble at room temperature. Once the solvent was replaced with acetonitrile, compound 1 crystallized perfectly.

Thus, through a manipulation with two solvents we isolated both high purity products on preparative scale and avoided time-consuming chromatography step [7,10,11,16]. The structures of 1 and 2 have been confirmed by spectroscopic methods. For compound 2, X-ray crystal structure was determined previously [17]. Our paper reports crystallographic data for Tröger's base 1 (Fig. 1).

## 2. Procedure

IR spectra were recorded on a Vertex 70 spectrometer. MS analysis was performed using Agilent 6890 N/5975 GC/MS instrument in the electron impact ionization mode at 70 eV.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded at 500 MHz and

125 MHz respectively on a Bruker Avance 500 spectrometer. Chemical shifts were referred to the signals of residual DMSO ( $\delta = 2.50$  ppm in  $^1\text{H}$  NMR and  $\delta = 39.50$  ppm in  $^{13}\text{C}$  NMR). Elemental analysis was performed using Thermo Scientific TM Flash 2000 CHNS/O analyser. Crystal structure was determined with Oxford Diffraction Xcalibur, Ruby, Gemini Ultra diffractometer.

**Synthesis and separation of compounds 1 and 2.** Ethyl *p*-aminobenzoate (40.5 g, 245 mmol) was added portionwise to mechanically stirred trifluoroacetic acid (400 mL), which was placed prior in an ice-salt bath and cold to  $-15$  °C. Once the addition was complete, the mixture was stirred for 15–20 min to lower slightly increased temperature. Whereupon paraformaldehyde (15.0 g, 500 mmol) was added portionwise, and stirring was continued for 16 h, meanwhile the mixture was gradually warmed to room temperature. The resulting mixture was evaporated in vacuum, the residue was treated consistently with crushed ice (500 g) and 25% ammonia solution (500 mL). The products were extracted with  $\text{CH}_2\text{Cl}_2$  ( $4 \times 250$  mL). Combined organic extracts were washed with saturated solution of  $\text{Na}_2\text{CO}_3$  (100 mL) and brine (100 mL), dried with  $\text{Na}_2\text{SO}_4$ , evaporated under reduced pressure. The residue was dissolved in refluxing toluene (200 mL), and transparent solution was left at room temperature. The solids formed were separated by filtration and washed with toluene ( $2 \times 50$  mL). Crystallization from toluene was repeated in the same manner giving pure compound **2** as pale yellow plates, yield 12.2 g (28%).

Mother liquor after first crystallization was concentrated in vacuum. The residue was twice recrystallized from of acetonitrile (120 and 130 mL, refrigerator) to afford pure Tröger's base **1** as colorless needles (15.9 g). Concentration and reprocessing of acetonitrile filtrates gave additional amounts of **1** (7.0 g). Total yield of **1**: 22.9 g (51%).

### 3. Data, value and validation

**2,8-Bis(ethoxycarbonyl)-6H,12H-5,11-methanodibenzo[b,f][1,5]diazocine 1.** Mp 155–156 °C (lit. [16] mp 152–153 °C). NMR and IR spectra were in agreement with reported [7,11,16].

**Ethyl 3-[4-(ethoxycarbonyl)phenyl]-3,4-dihydroquinazoline-6-carboxylate 2.** Mp 185–187 °C (lit. [11] mp 190–191 °C), IR (film),  $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ : 3388, 3076, 2980, 2904, 1691, 1631, 1591, 1558.  $^1\text{H}$  NMR spectrum (500 MHz,  $\text{DMSO}-d_6$ ),  $\delta$ , ppm (*J*, Hz): 8.01 (2H, br d, *J* = 8.9, Ar); 7.95 (1H, s,  $\text{CH}=\text{N}$ ); 7.81 (1H, dd, *J* = 8.2, *J* = 1.8, Ar); 7.75 (1H, br d, *J* = 1.8, Ar); 7.47 (2H, br d, *J* = 8.9, Ar); 7.16 (1H, d, *J* = 8.2, Ar); 5.06 (2H, s,  $\text{ArCH}_2$ ); 4.31 (2H, q, *J* = 7.1,  $\text{CH}_3\text{CH}_2\text{O}$ ); 4.29 (2H, q, *J* = 7.1,  $\text{CH}_3\text{CH}_2\text{O}$ ); 1.33 (3H, t, *J* = 7.0,  $\text{CH}_3$ ), 1.31 (3H, t, *J* = 7.0,  $\text{CH}_3$ ).  $^{13}\text{C}$  NMR spectrum (125 MHz,  $\text{DMSO}-d_6$ ),  $\delta$ , ppm: 165.24 (C); 165.20 (C); 147.6 (CH); 145.8 (C); 145.1 (C); 130.5 (2CH); 129.6 (CH); 127.4 (CH); 126.5 (C); 125.0 (C); 124.2 (CH); 122.3 (C); 116.9 (2CH); 60.63 ( $\text{CH}_2$ ); 60.58 ( $\text{CH}_2$ ); 45.5 ( $\text{CH}_2$ ); 14.2 (2 $\text{CH}_3$ ). NMR spectra were similar to reported [11]. MS (ESI), *m/z*: 353.2 [*M* + *H*] $^+$ .

**Crystal data for 1.**  $\text{C}_{21}\text{H}_{22}\text{N}_2\text{O}_4$  (*M* 366.41), orthorhombic, *Pca*2 $_1$ , *a* = 10.2541(5), *b* = 22.8519(11) and *c* = 8.0645(3) Å, *V* = 1889.72(15) Å $^3$ , *Z* = 4,  $d_{\text{calc}}$  = 1.288 g  $\text{cm}^{-3}$ , *F*(000) = 776. Single crystal (colorless needle, dimensions 0.029 × 0.067 × 0.65 mm) was selected and intensities of 5472 reflections were measured with Oxford Diffraction Xcalibur, Ruby, Gemini Ultra diffractometer at 295 K [ $\omega$  scans,  $\lambda(\text{Cu K}\alpha)$  = 1.54184 Å,  $\mu$  = 0.733  $\text{mm}^{-1}$ ,  $2\theta_{\text{max}}$  = 134.456°]. Final *R* factors: *R* $_1$  = 0.0532 [2253 reflections with *I* > 2σ(*I*)], *wR* $^2$  = 0.1589 (all reflections), GOF = 1.030. The structure was solved with the SHELXT [18] and refined with the SHELXL-2016/6 program [19]. Molecular graphics were prepared using Olex2 program [20].

CCDC 1934607 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk>.

The supplementary materials file containing copies of NMR spectra is available from the journal website.

### Declaration of Competing Interest

None.

### Acknowledgments

This work was supported by the Belarusian Republican Foundation for Basic Research (Grant Numbers: X18KOR-003; X17INDG-006). We thank the PC $^2$  technological platform of UNamur for access to single-crystal X-ray diffraction instrument.

### Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.cdc.2020.100339.

### References

- [1] J. Tröger, Ueber einige mittelst nascirenden Formaldehydes entstehende Basen, *J. Prakt. Chem.* 36 (1887) 225–245.
- [2] B. Dolensky, J. Elguero, V. Kral, C. Pardo, M. Valik, Current Tröger's base chemistry, *Adv. Heterocycl. Chem.* 93 (2007) 1–56.
- [3] S. Sergeev, Recent developments in synthetic chemistry, chiral separations, and applications of Tröger's base analogues, *Helv. Chim. Acta* 92 (2009) 415–444.
- [4] O.V. Runarsson, J. Artacho, K. Warnmark, The 125th anniversary of the Tröger's base molecule: synthesis and applications of Tröger's base analogues, *Eur. J. Org. Chem.* (2012) 7015–7041.

- [5] R.G. Kostyanovsky, V.R. Kostyanovsky, G.K. Kadorkina, K.A. Lyssenko, Asymmetric three-coordinated nitrogen compounds: spontaneous resolution and absolute asymmetric synthesis, *Mendeleev Commun.* 13 (2003) 111–113.
- [6] C.S. Hawes, C.M. Fitchett, S.R. Batten, P.E. Kruger, Synthesis and structural characterisation of a Co(II) coordination polymer incorporating a novel dicarboxy-Tröger's base/bis-pyrazole mixed ligand system, *Inorg Chim. Acta* 389 (2012) 112–117.
- [7] S. Goswami, K. Ghosh, S. Dasgupta, Tröger's base molecular scaffolds in dicarboxylic acid recognition, *J. Org. Chem.* 65 (2000) 1907–1914.
- [8] S. Goswami, K. Ghosh, Molecular recognition: chain length selectivity studies of dicarboxylic acids by the cavity of a new Tröger's base receptor, *Tetrahedron Lett.* 38 (1997) 4503–4506.
- [9] J. Jensen, K. Wannmark, Synthesis of halogen substituted analogues of Tröger's base, *Synthesis* (2001) 1873–1877.
- [10] D. Didier, B. Tylleman, N. Lambert, C.M.L. Vande Velde, F. Blockhuys, A. Collas, S. Sergeyev, Functionalized analogues of Tröger's base: scope and limitations of a general synthetic procedure and facile, predictable method for the separation of enantiomers, *Tetrahedron* 64 (2008) 6252–6262.
- [11] M.D.H. Bhuiyan, K.-X. Zhu, P. Jensen, A.C. Try, Synthesis of symmetric diester-functionalised Tröger's base analogues, *Eur. J. Org. Chem.* (2010) 4662–4670.
- [12] Mechanistic explanation and discussion see in ref. [4].
- [13] N. Gruber, J.E. Diaz, L.R. Orelli, Synthesis of dihydroquinazolines from 2-aminobenzylamine:  $N^3$ -aryl derivatives with electron-withdrawing groups, *Beilstein J. Org. Chem.* 14 (2018) 2510–2519.
- [14] C. Wu, J. Wang, X.-Y. Zhang, G.-K. Jia, Z. Cao, Z. Tang, X. Yu, X. Xub, W.-M. He, Palladium-catalyzed selective synthesis of 3,4-dihydroquinazolines from electron-rich arylamines, electron-poor arylamines and glyoxalates, *Org. Biomol. Chem.* 16 (2018) 5050–5054.
- [15] Q. Chen, Z. Mao, K. Gao, F. Guo, L. Sheng, Z. Chen, Synthesis of 1,2-dihydroquinazolines via rearrangement of indazolium salts, *J. Org. Chem.* 83 (2018) 8750–8758.
- [16] Z. Li, X. Xu, Y. Peng, Z. Jiang, C. Ding, X. Qian, An unusual synthesis of Tröger's bases using DMSO/HCl as formaldehyde equivalent, *Synthesis* (2005) 1228–1230.
- [17] M.D.H. Bhuiyan, J.K. Clegg, A.C. Try, Ethyl 3-[4-(ethoxycarbonyl)phenyl]-3,4-dihydroquinazoline-6-carboxylate, *Acta Crystallogr. E* 63 (2007) o308–o310.
- [18] G.M. Sheldrick, SHELXT – Integrated space-group and crystal-structure determination, *Acta Crystallogr. A* 71 (2015) 3–8.
- [19] G.M. Sheldrick, Crystal structure refinement with SHELXL, *Acta Crystallogr. C* 71 (2015) 3–8.
- [20] O.V. Dolomanov, L.J. Bourhis, R.J. Gildea, J.A.K. Howard, H.J. Puschmann, OLEX2: a complete structure solution, refinement and analysis program, *J. Appl. Crystallogr.* 42 (2009) 339–341.